


Patient Name : MR SHAIKH RASHEED NABI AHMED
Age and Gender : 45 / Years / M
Aadhaar Card : 747968902609
Patient Location : Mumbai
Patient Contact No. : 7079283336
Referring Doctor : SELF
Centre : COOPER HOSPITAL

Sample Collection Time : 04.03.2022 11:00 PM
Sample Accepted On : 05.03.2022 01:05 AM
Reported On : 05.03.2022 04:36 AM
Sample UID No. 
01DSK001922
Organization- : F South Mumbai

SARS CoV -2 REAL TIME PCR (COVID-19)

Test Done	Observed Value
-----------	----------------

SARS CoV -2 REAL TIME PCR (COVID-19)

SPECIMEN

SARS-CoV-2 RNA

Method

Note :

Marked parameters are not in the scope of NABL accreditation.

Kit Specification - Qiagen Covid-19 One Step RT-PCR kit.

Interpretation(s)

SARS CoV -2 REAL TIME PCR-

SARS-CoV-2, formerly known as 2019-nCoV, is the causative agent of the coronavirus disease 2019 (COVID-19). Main symptoms of the disease include fever, cough and shortness of breath. The virus is spread via person-to-person contact through respiratory droplets produced when a person coughs or sneezes. The SARS-CoV-2 RNA is generally detectable in nasopharyngeal/oropharyngeal swabs during the acute phase of infection. Positive results are indicative of active infection. Real Time PCR assay targets specific genes and can be used for diagnosis of SARS-CoV-2 virus infection which contributes to severe upper respiratory distress and complications.

Positive result indicates that RNA from SARS-CoV-2 was detected in the specimen, and the patient is considered infected with the virus and presumed to be contagious. Negative test result for this test means that SARS-CoV-2 RNA was not detected in the specimen above the limit of detection of the assay.


Limitations:

- Negative results do not preclude COVID-19 and should not be used as the sole basis for patient management decisions. Negative results must be combined with clinical observations, patient history, and epidemiological information.
- Positive results do not rule out bacterial infection or co-infection with other viruses.
- Optimum specimen types and timing for peak viral levels during infections caused by 2019-nCoV have not been determined. Collection of multiple specimens (types and time points) from the same patient may be necessary to detect the virus.
- Follow-up testing may particularly be important if patient has a clinical picture of viral pneumonia, a potential exposure history, and/or radiographic findings (chest CT or MRI scan) consistent with COVID -19 pneumonia. However repeat testing in the near-term after clearance (within 90 days) should be avoided as prolonged shedding of non-viable virus is not uncommon.
- Ct values generated from different assay systems within the same laboratory, or from different laboratories, are not directly comparable and do not

~~~ END OF REPORT ~~~




Dr Sudhir D. Khodwekar  
PHD Biotechnology



Dr Minal Kusalkar  
MD Microbiologist

Patient Name : MR SHAIKH RASHEED NABI AHMED  
Age and Gender : 45 / Years / M  
Aadhaar Card : 747968902609  
Patient Location : Mumbai  
Patient Contact No. : 7079283336  
Referring Doctor : SELF  
Centre : COOPER HOSPITAL

Sample Collection Time : 04.03.2022 11:00 PM  
Sample Accepted On : 05.03.2022 01:05 AM  
Reported On : 05.03.2022 04:36 AM  
Sample UID No.   
Organization- : F South Mumbai

**SARS CoV -2 REAL TIME PCR ( COVID-19 )**

| Test Done | Observed Value |
|-----------|----------------|
|-----------|----------------|

**SARS CoV -2 REAL TIME PCR ( COVID-19 )**

necessarily reflect the same viral load due to inter-assay and inter-laboratory variability.

- Variation in timing of sample collection, fluctuations in virus shedding, and difference between detection limit of different testing methods within same or different labs could lead to variation in results particularly during initial phase of infection.
- If the virus mutates in the RT-PCR target region, 2019-nCoV may not be detected or may be detected less predictably. Inhibitors or other types of interference may produce a false negative result.
- The performance of this test has not been established for monitoring treatment of 2019-nCoV infection.

**Note:**

Test is performed using ICMR approved Kit.

**References:**

1. Laboratory testing for coronavirus disease 2019 (COVID-19) in suspected human cases. Interim guidance. World Health Organization.
2. Druce et al. JCM. 2011
3. N. Engl. J. Med. 2020, 382, 929–936

ICMR Registration no - MEDCMMH

NABL Certificate no - MC-4413

Results Entered By : E10023


Results Approved By : Dr Minal Kusalkar

SAMPLE PROCESSED AT - MEDILAB DIAGNOSTIC CENTER SEWREE

~~~ END OF REPORT ~~~



Dr Sudhir D. Khodwekar
PHD Biotechnology



Dr Minal Kusalkar
MD Microbiologist